



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$= L(1+x^2) - \frac{1}{x^2} L(1+x^2) + 1 + \frac{4}{x} (\tan^{-1}x - x).$$

$$\therefore S(+1) = 4\left(\frac{\pi}{4} - 1\right) + 1 = \frac{1}{3 \cdot 1 \cdot 2} - \frac{1}{5 \cdot 2 \cdot 3} + \dots$$

$$\therefore \pi = 3 + \frac{1}{3} \cdot \frac{1}{1 \cdot 2} - \frac{1}{5} \cdot \frac{1}{2 \cdot 3} + \dots$$

Also solved by G. B. M. Zerr, and V. M. Spunar.

339. Proposed by E. B. ESCOTT, University of Michigan, Ann Arbor, Mich.

Prove that if $a_1 < 2$ and $a_n = a_{n-1}^2 - 2$, $\frac{1}{a_1} + \frac{1}{a_1 a_2} + \frac{1}{a_1 a_2 a_3} + \dots$
 $= \frac{1}{2} [a_1 - \sqrt{(a_1^2 - 4)}].$

Solution by the PROPOSER.

Let the roots of the equation $x^2 - a_1 x + 1 = 0$ be a and $1/a$. Then

$$\begin{aligned} a_1 &= a + \frac{1}{a}, \\ a_2 &= a^2 + \frac{1}{a^2}, \\ a_3 &= a^4 + \frac{1}{a^4}, \\ &\dots \end{aligned}$$

The series becomes

$$\frac{1}{a + \frac{1}{a}} + \frac{1}{\left(a + \frac{1}{a}\right)\left(a^2 + \frac{1}{a^2}\right)} + \dots + \left(a - \frac{1}{a}\right) \left[\frac{1}{a^2 - \frac{1}{a^2}} + \frac{1}{a^4 - \frac{1}{a^4}} + \frac{1}{a^8 - \frac{1}{a^8}} \dots \right]$$

$$\text{Since } \frac{1}{a^2 - \frac{1}{a^2}} = \frac{1}{2} \left(\frac{a + \frac{1}{a}}{a - \frac{1}{a}} - \frac{a^2 + \frac{1}{a^2}}{a^2 - \frac{1}{a^2}} \right),$$

each term of the series may be written as the difference of two fractions, *i. e.*

$$\frac{a - \frac{1}{a}}{2} \left[\left(\frac{a + \frac{1}{a}}{a - \frac{1}{a}} - \frac{a^2 + \frac{1}{a^2}}{a^2 - \frac{1}{a^2}} \right) + \left(\frac{a^2 + \frac{1}{a^2}}{a^2 - \frac{1}{a^2}} - \frac{a^4 + \frac{1}{a^4}}{a^4 - \frac{1}{a^4}} \right) + \dots \right]$$

$$= -\frac{1}{2} \left[\frac{a + \frac{1}{a}}{a - \frac{1}{a}} - \frac{a^{2^n} + a^{-2^n}}{a^{2^n} - a^{-2^n}} \right].$$

When $a > 1$, the limit of this sum is

$$-\frac{1}{2} \left(\frac{a + \frac{1}{a}}{a - \frac{1}{a}} - 1 \right) = \frac{1}{a}.$$

Therefore, the sum of the series is the smaller root of the equation $x^2 - a_1x + 1 = 0$, viz., $\frac{a_1 - \sqrt{a_1^2 - 4}}{2}$.

NOTE.—This furnishes a very rapid method for finding square roots to a considerable number of decimals. Example.—Let $a = 16$,

$$8 - 3\sqrt{7} = \frac{1}{16} + \frac{1}{16.254} + \frac{1}{16.254.645.64514} + \dots$$

and three terms of the series give $\sqrt{7}$ to 18 decimals, four terms will give 37 decimals, etc.

Also solved by S. Lefschetz.

GEOMETRY.

365. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

Given the coordinates of the four vertices of the tetrahedron, (x_1, y_1, z_1) ; (x_2, y_2, z_2) ; (x_3, y_3, z_3) ; (x_4, y_4, z_4) : find volume and express it by a determinant.

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa., and I. W. SMITH, A. M., Assistant Professor of Mathematics, North Dakota Agricultural College.

Let Δ = area of triangular base BCD of the tetrahedron, p the perpendicular from the vertex A on the base BCD , V its volume, $(x - x_2) \cos \alpha + (y - y_2) \cos \beta + (z - z_2) \cos \gamma = 0$ the equation to the plane of BCD .

Then $\Delta \cos \gamma$ is the projection of the area BCD on the plane xy , and (x_2, y_2) , (x_3, y_3) , (x_4, y_4) are its angular points.